



The interpreter's brain during rest - Hyperconnectivity in the frontal lobe

Klein, Carina ; Metz, Silvana Iris ; Elmer, Stefan ; Jäncke, Lutz

Abstract: Language in its highest complexity is a unique human faculty with simultaneous translation being among the most demanding language task involving both linguistic and executive functions. In this context, bilingually grown up individuals as well as simultaneous interpreters (SIs) represent appropriate groups for studying expertise-related neural adaptations in the human brain. The present study was performed to examine if a domain-specific neural network activation pattern, constituted by brain regions involved in speech processing as well as cognitive control mechanisms can be detected during a task-free resting state condition. To investigate this, electroencephalographic (EEG) data were recorded from 16 SIs and 16 age and gender-matched multilingual control subjects. Graph-theoretical network analyses revealed interhemispheric hyperconnectivity between the ventral part of the prefrontal cortex (pars opercularis and pars triangularis) and the dorsolateral prefrontal cortex (DLPFC) in language experts compared to multilingual controls in the alpha frequency range. This finding suggests that the high cognitive demands placed on simultaneous interpreting lead to an increased neural communication between prefrontal brain regions essentially engaged in supporting executive control-a neural fingerprint that is even detectable during rest.

DOI: <https://doi.org/10.1371/journal.pone.0202600>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-157017>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Klein, Carina; Metz, Silvana Iris; Elmer, Stefan; Jäncke, Lutz (2018). The interpreter's brain during rest - Hyperconnectivity in the frontal lobe. PLoS ONE, 13(8):e0202600.

DOI: <https://doi.org/10.1371/journal.pone.0202600>


RESEARCH ARTICLE

The interpreter's brain during rest — Hyperconnectivity in the frontal lobe

Carina Klein^{1*}, Silvana Iris Metz¹, Stefan Elmer¹, Lutz Jäncke^{1,2,3,4,5}

1 Division of Neuropsychology, Department of Psychology, University of Zurich, Zurich, Switzerland, **2** International Normal Aging and Plasticity Imaging Center (INAPIC), University of Zurich, Zurich, Switzerland, **3** Center for Integrative Human Physiology (ZIHP), University of Zurich, Zurich, Switzerland, **4** University Research Priority Program (URPP), Dynamic of Healthy Aging, University of Zurich, Zurich, Switzerland, **5** Department of Special Education, King Abdulaziz University, Jeddah, Saudi Arabia

* carina.klein@uzh.ch



Abstract

Language in its highest complexity is a unique human faculty with simultaneous translation being among the most demanding language task involving both linguistic and executive functions. In this context, bilingually grown up individuals as well as simultaneous interpreters (SIs) represent appropriate groups for studying expertise-related neural adaptations in the human brain. The present study was performed to examine if a domain-specific neural network activation pattern, constituted by brain regions involved in speech processing as well as cognitive control mechanisms can be detected during a task-free resting state condition. To investigate this, electroencephalographic (EEG) data were recorded from 16 SIs and 16 age and gender-matched multilingual control subjects. Graph-theoretical network analyses revealed interhemispheric hyperconnectivity between the ventral part of the prefrontal cortex (pars opercularis and pars triangularis) and the dorsolateral prefrontal cortex (DLPFC) in language experts compared to multilingual controls in the alpha frequency range. This finding suggests that the high cognitive demands placed on simultaneous interpreting lead to an increased neural communication between prefrontal brain regions essentially engaged in supporting executive control—a neural fingerprint that is even detectable during rest.

OPEN ACCESS

Citation: Klein C, Metz SI, Elmer S, Jäncke L (2018) The interpreter's brain during rest — Hyperconnectivity in the frontal lobe. PLoS ONE 13(8): e0202600. <https://doi.org/10.1371/journal.pone.0202600>

Editor: Yuka Kotozaki, Iwate Medical University, JAPAN

Received: June 19, 2018

Accepted: August 6, 2018

Published: August 23, 2018

Copyright: © 2018 Klein et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Funding: This research was supported by the Swiss National Science Foundation (SNSF, Grant No 320030B_138668 granted to Lutz Jäncke. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Simultaneous interpreting is one of the most complex language tasks involving various linguistic and cognitive functions. In fact, it requires a continuous encoding of the input language, the maintenance of the heard information in verbal short-term memory while at the same time translating the information into the output language, inhibiting the articulation of the source, and producing the target language [1, 2]. In the last two decades, first magnetic resonance imaging (MRI) and positron emission tomography (PET) studies have addressed long-term training effects of simultaneous interpreting on the structural [3–5] and the functional [6, 7] architecture of the human brain. Interestingly, results of structural MRI studies revealed differences in brain regions which are also known to be altered in bilinguals. For example,

reduced gray matter volume was observed in the middle anterior cingulate gyrus, a brain region which is involved in attentional functions supporting conflict monitoring and error detection [8–10]. Furthermore, gray matter volume was altered in the pars triangularis and its right hemispheric homologue, brain regions playing a role in syntactic processing [11], prosody perception [12], language switching, attention and working memory [13–15]. Also, the left supramarginal gyrus contributing to phonetic processing [16, 17] as well as the brain regions regulating speech production and articulation, i.e. left middle anterior insula [18, 19], caudate nucleus [20] and left pars opercularis [3, 11, 20] were found to have reduced morphological measures. Decreased fractional anisotropy values were found in brain areas regulating speech articulation and sensory-to-motor coupling mechanisms (left anterior insula, upper part of the corticospinal tract, and dorsal part of the right caudate nucleus) as well as controlling interhemispheric information transfer (genu and splenium of the corpus callosum) [4]. A longitudinal MRI study in which subjects were scanned before and after a 15-month period of simultaneous interpreting training showed an increase of cortical thickness over time in brain regions involved in phonetic processing (left posterior superior temporal gyrus, anterior supramarginal gyrus and planum temporale), propositional speech (right angular gyrus), working memory (right dorsal premotor cortex) as well as in executive control functions and attention (right parietal lobule) [5].

In a functional MRI study by Hervais-Adelman and colleagues [7], neural activity during simultaneous interpreting was examined pre- and post-training in contrast to speech repetition (shadowing). The authors found a reduced training-related recruitment in the caudate nucleus, interpreting their findings as a decreased demand on multilingual language control as the task becomes more automatized over the course of an intense training. In 2000, Rinne and colleagues measured brain activation in SIs in a PET scanner during different directions of interpreting, i.e. interpreting into the native language (from L2 to L1), into the non-native language (from L1 to L2—the cognitive more demanding condition) and during shadowing (simultaneous aloud repetition of heard text) [6]. The following contrasts were built: 1) L1 to L2 translation versus shadowing L1, 2) L2 to L1 translation versus shadowing L2 as well as 3) L1 to L2 translation minus shadowing L1 versus L2 to L1 translation minus shadowing L2. Across all contrasts, the authors found mainly left-lateralized effects of simultaneous interpreting: the first condition elicited neural activity in the left frontal lobe (Brodmann area (BA) 6 and BA46). The second condition showed involvement of the left frontal lobe (BA6, BA45), the left inferior temporal zone (BA20/28) as well as the right cerebellum. Contrasting the two interpreting directions (third condition) revealed an increased activation in BA44. The authors showed that cerebral activation patterns vary according to the interpreting direction, recruiting more regions when interpreting in the more effortful direction. Thereby, brain regions are activated that are involved in verbal encoding, processing semantic information, working memory, and when solving effortful tasks.

To the best of our knowledge, only one fMRI study exists that examined spontaneous neural activity during a task-free resting state condition in different types of interpreters: SIs, consecutive interpreters and translators [21]. The authors of this study found hyperconnectivity in SIs compared to the other groups between the left frontal pole and the left middle temporal gyrus as well as between the former region and the left pars opercularis and triangularis.

Despite previous insight into differential brain activation during simultaneous interpreting, to date nothing is known about neural activation patterns during resting state in SIs compared to a multilingual control group. In the context of long-term training and its influence on neural oscillatory activity during resting state, it has repeatedly been shown that repetitive task-specific activations lead to altered network patterns during rest across a variety of domains. For example, a resting state EEG study in professional string players reported increased intra-

and interhemispheric functional connectivity between brain regions that are typically involved in music processing and production, such as somatosensory, auditory, and prefrontal regions as well as Broca's area [22]. Expertise-related findings have also been reported in professional chess [23] or badminton players [24] and dancers [25]. These studies emphasize the recruitment of expertise-characteristic brain regions during a resting state period even when not being confronted with the respective stimuli or situation at the moment of data collection.

Thus, instead of examining expertise-specific activation patterns during a task condition, the aim of the present study was to find a putative modulation of resting state network characteristics as a function of long-term simultaneous interpreting training. In this context, we expect to find altered functional connectivity measures in SIs compared to healthy age- and gender matched multilingual control subjects between brain regions that are involved in language processing and control as well as in administering cognitive control functions, such as Broca's area [3, 11], regions in the ventral and dorsal prefrontal cortex and the posterior part of the middle temporal gyrus and the temporal pole as well as the inferior parietal lobe [5, 6]. Examining EEG-based connectivity measures in the source space allows us to draw conclusions on the underlying brain structures constituting an altered functional network architecture in the SIs compared to the control group. The use of EEG in contrast to fMRI further bears the advantage to disentangle network characteristics in separate frequency ranges. This provides further information on a functionally relevant level than simply discussing the involved brain regions per se. Here, we particularly focused on two frequency ranges being involved in executive control functions such as working memory (i.e. the theta frequency band) [26], inhibition and attention (i.e. the alpha frequency band) [27–29].

Materials and methods

Subjects

Sixteen female right-handed SIs, all graduates/students from a local Master's in Applied Linguistics program on simultaneous interpreting (mean age = 34.7, standard deviation (SD) = 9.4 years; age of acquisition of the second language (AoA L2) = 9.4, SD = 3.4; mean age of commencement (AoC) = 27.0, SD = 3.5 years; mean years of experience (YoE) = 7.7, SD = 8.3 years; estimated number of cumulated training hours during life = 4298.1 hours, SD = 6038.9 hours) and sixteen multilingual control subjects (mean age = 34.3, SD = 9.0 years; AoA L2 = 9.9, SD = 3.8) participated in this study. Subjects were matched for handedness [30], age, gender and multilingualism (for detailed information see S1 and S2 Tables). One subject per group grew up bilingually. Language experience was measured with an in-house questionnaire collecting information about mother tongue, bilingualism, foreign languages and the respective AoCs, information about simultaneous interpreting activity, usage of languages (speaking, movie watching, reading, working environment, et cetera) as well as self-reported language proficiency in speaking and writing in everyday communications on a scale from 1 (basic) to 5 (fluent; see S1 File). This questionnaire was used in previous studies [3, 31, 32]. Participants with professional music education were excluded from the study. Musical aptitude was measured with the Advanced Measures of Music Audition (AMMA) test [33]. In this test, participants listen to 30 trials consisting of two successive short piano melodies each. After every single trial, the subject has to decide if the two melodies are either identical or differ in rhythmic or tonal specification from each other. Furthermore, an in-house questionnaire on musical history was applied. In this questionnaire, which was also used in previous studies (see for example [31, 34, 35]), the played musical instruments, musical education as well as training hours per day and week in specific age ranges are indicated (see S2 File). None of the participants reported any current or past neurological, psychiatric, or neuropsychological disease, nor medication or drug abuse.

Written informed consent was obtained from all participants and they were paid for participation. The study was approved by the local ethics committee (Kantonale Ethikkommission of the canton of Zurich) according to the declaration of Helsinki.

Cognitive capability

For testing cognitive capability, participants underwent the KAI (Kurztest für allgemeine Basisgrösse der Informationsverarbeitung) [36] examining the actual cognitive capability (fluid intelligence) based on short-term memory capacity and speed of information processing as well as the MWT-B (Mehrfachwahl-Wortschatz Intelligenztest) [37] testing for crystalline intelligence. In the KAI test, the time for reading aloud four random sequences of visually presented letters as well as the capacity of how many items (numbers and letters, one up to nine items) can be remembered and orally repeated is measured. In the MWT-B, subjects have to choose 37 times the "real" word out of six pseudowords.

Experimental procedure

Before the EEG recording, participants were tested for their cognitive capabilities and filled out the handedness questionnaire as well as in-house questionnaires about health, music and language skill. After electrode application, participants were placed in a light-dimmed and sound shielded Faraday cage. They sat down in a comfortable chair at a table with a 19-inch monitor at a distance of approximately 75 cm. Subjects were asked to sit calm and relaxed and to avoid strong movements during recording. The EEG paradigm consisted of a task-free period of 3 min eyes open (black fixation cross on a white screen) and 3 min eyes closed (black screen). Afterwards, a verbal Sternberg paradigm was presented according to a previous study by Klein et al. 2016 [38] as well as a post-task resting state period. The present study only covers data analysis of the pre-task resting state condition (3 min eyes open and 3 min eyes closed separately). Presentation of the stimulus material was controlled with the Presentation software (Neurobehavioral Systems, Berkeley, USA, <http://www.neurobs.com>).

EEG recording and processing

Continuous EEG was recorded with a 32-channel cap (EASYCAP GmbH, Herrsching, Germany) with a sampling rate of 500 Hz and a high-pass butterworth filter with zero phase distribution (0.1 Hz) by using the BrainVision Recorder Software (Brainproducts, Gilching, Germany, <http://www.brainproducts.com>). The electrodes (silver/silver chloride sintered ring electrodes) were located at frontal, temporal, parietal and occipital scalp sites according to the international 10–20 system (Fp1, Fp2, F7, F3, Fz, F4, F8, FT7, FC3, FCz, FC4, FT8, T7, C3, Cz, C4, T8, TP9, TP7, CP3, CPz, CP4, TP8, TP10, P7, P3, Pz, P4, P8, O1, Oz, O2). The nose electrode was used as online reference. Electrode impedances were kept below 10 k Ω by applying abrasive electrically conductive gel.

Preprocessing of the data was performed offline by using the BrainVision Analyzer software (version 2.1, Brainproducts, Gilching, Germany, <http://www.brainproducts.com>). Data were band pass filtered between 0.1–100 Hz (high- and low-pass butterworth filter with zero phase distribution), including a notch filter (50 Hz). An independent component analysis was applied to remove eye blinks and eye movement artifacts [39]. Remaining artifacts were removed by a semi-automatic raw data inspection (maximal allowed voltage steps: 50 μ V, maximal allowed differences of values: 200 μ V over 200 ms, allowed amplitude range: -200 to 200 μ V, lowest allowed activity in intervals: 0.5 μ V over 100 ms) and bad channels were interpolated. After artifact cleaning, data were again band pass filtered between 1 Hz and 40 Hz (high- and low-pass butterworth filter with zero phase distribution). The 3 min eyes open and

eyes closed resting state periods were segmented into 2 s segments each. Only artifact free segments entered connectivity analysis (a minimum of 74 two second segments per subject; mean number of segments: 88, SD = 2.8).

Connectivity analysis in the source space. Functional connectivity is a measure displaying statistical dependencies of neural activity between different regions of interest (ROIs) [40]. In graph-theoretical approaches, a node represents the ROI, while an edge refers to any kind of connection value between the ROIs. As suggested by literature, connectivity analysis in the source space represents a methodological advantage over measuring coherences on the scalp level (between electrodes) and provides simplified result interpretation [41, 42]. Although there is some argument about the reliability of source reconstruction of EEG data, literature exists across diverse conditions [43–45] besides cross-validation studies with functional MRI (fMRI) [46, 47] or PET [48] that show meaningful results for the source reconstruction method used in the present study [49]. Connectivity analyses were performed using the Low Resolution Electromagnetic Tomography (LORETA, Standardized & Exact) KEY toolbox (version 2016–05; <http://www.uzh.ch/keyinst/loreta.htm>) between all 84 BAs (42 in each hemisphere) implemented in the LORETA KEY software [49–51]. In LORETA, electrically active neuronal generators are calculated based on the recorded electric scalp potentials, which are transformed into current density values (A/cm^2) for every voxel, assuming similar activation among neighbored voxels. Computations are made in a realistic gray-matter head model using the 152MNI (Montreal Neurological Institute) template by taking into account the effect of volume conduction. In LORETA, the intracerebral space is partitioned into 6239 voxels with a spatial resolution of 5 mm. Because of the assumption of smoothness of data (i.e. neighbored voxels show highly correlated activity), connectivity (here linear instantaneous dependency values, a measure of zero-phase, zero-lag covariances) was calculated between the centroid voxels of the 84 BAs [52]. Linear instantaneous dependencies are a validated measure for calculating distributed networks between predefined brain regions and were repeatedly shown to be affected by long- and short-term training, i.e. in the frame of expertise research [22, 53–55]. The ROI-based transformation matrix was created using the automatic approach. The resulting transformation matrix with a signal-to-noise ratio of 1 was used to calculate instantaneous connectivity values in the theta (4–7 Hz), lower (8–10 Hz) and upper alpha (10.5 Hz–12 Hz) frequency ranges.

As a result, LORETA provides connectivity matrices with linear instantaneous coherence values, one value for every single subject in every single frequency range between all BAs across the 3 min resting state periods. For better result interpretation, the Harvard-Oxford and the Juelich-Histological cortical atlases (<http://fsl.fmrib.ox.ac.uk/fsl/fslwiki/Atlases>) implemented in the FSL software (<http://fsl.fmrib.ox.ac.uk/fsl/fslwiki/FSL>) were used for a detailed assignment of the MNI coordinates of the centroid voxels of the ROIs (BAs) to the hyper-connected brain regions, respectively.

Network-based statistic

For evaluating between-group differences in functional connectivity, the individual connectivity matrices obtained from the LORETA KEY software were subjected to network-based statistic (NBS), separately for the frequency bands of interest [56]. In traditional network analyses, t-tests for each connection of the connectivity matrix are performed and corrected for multiple comparisons without considering dependencies between the connections building a (sub)network. In contrast, NBS tests the network as a whole and is based on a non-parametric supra-threshold cluster test as often applied in fMRI analyses [57]. On the basis of a general linear model approach, the t-test module in NBS was used to compare the individual connectivity

matrices between the two groups. By using the component extent option, the extent of functional connections comprising the contrast or effect of interest (i.e., the group difference of the functional connectome during resting state) is examined. For finding a subnetwork in which the two groups differ, sensitivity (set) thresholds are chosen in advance (here t -thresholds) to define which edges form a supra-threshold network. The t -threshold must be chosen arbitrarily in an explorative way. However, as these t -thresholds do not represent the actual alpha error probabilities, this procedure does not affect the false-positive rate of the actual permutation testing. The size, that is the number of edges of the biggest supra-threshold subnetwork, also called the biggest component is then used for test statistics. We controlled the alpha error probability ($p < 0.05$) by permuting 10000 times the group labels of the subjects and the size of the biggest component. A family-wise error (FWE)-corrected p -value is estimated by counting the number of permutations for which the size of the largest random supra-threshold network is bigger than the one of the real data and dividing it by the number of performed permutations [56]. Here, the connectivity matrices were tested for a range of set thresholds between $t = 2.0$ to 4.0 in increments of 0.1 . Here, we report the biggest obtained subnetworks for $t = 3.3$. P -values for the network-based measures are one-tailed since a directed contrast is tested. The obtained subnetworks were visualized with the BrainNet Viewer (version 1.53; <http://www.nitrc.org/projects/bnv/>) [58].

Statistics of behavioral data and correlation analyses

Demographical data and cognitive capability values were analyzed in the IBM software SPSS Statistics 25 (SPSS Inc, Chicago, IL, USA; <https://www.ibm.com/products/spss-statistics>). To investigate group comparisons of age and cognitive capability, age and the KAI score were compared with the Mann-Whitney U test (according to deviations from a normal distribution) and a two-tailed unpaired t -test was performed comparing MWT-B values.

To test for brain-behavior relationships, we calculated correlation analyses according to Spearman's rho (two-tailed) between the mean connectivity value of the entire network and the language characteristics YoE, hours of training per week and life in simultaneous interpreting, and AoC.

Results

Autobiographical and behavioral data

The two groups did not differ in age ($t_{(30)} = 0.115$, $p = 0.909$, two-tailed), AoA L2 ($z = -0.710$, $p = .478$) or in performance in the KAI test ($z = -1.543$, $p = 0.123$, two-tailed). However, SIs reached a significant higher score in the MWT-B test ($t_{(30)} = 2.66$, $p = 0.012$, two-tailed). This might be explained by the fact that the MWT-B relies very much on language experience. Although the control subjects were matched in the number of learned languages, a professional SI is still much more experienced and thus outperforms untrained controls.

Network-based statistic

[Table 1](#) contains the assignments of the centroid voxels of the BAs constituting the significant network obtained from the NBS analysis to MNI coordinates.

The SIs showed increased functional connectivity values only in the lower alpha frequency band for a t -value of 3.3 compared to the control group (SIs: group mean of the connectivity values = 5.42 , $SD = 0.46$; controls: group mean of the connectivity values = 3.78 , $SD = 0.37$; $p = 0.046$, FWE corrected; effect size $r = 0.56$ according Cohen's $d \geq 1.35$). In particular, the network consisted of eight nodes and eight edges between the left DLPFC, left pars opercularis

Table 1. Specification of Brodmann areas.

BA	MNI coordinates (x, y, z)	Brain region
BA9	(-30, 30, 35)	middle frontal gyrus (dorsolateral prefrontal cortex)
BA9	(30, 30, 35)	middle frontal gyrus (dorsolateral prefrontal cortex)
BA44	(-50, 10, 15)	Broca's area (pars opercularis)
BA44	(55, 10, 15)	right homologue of Broca's area (pars opercularis)
BA45	(-50, 20, 15)	Broca's area (pars triangularis)
BA45	(50, 20, 15)	right homologue of Broca's area (pars triangularis)
BA46	(-45, 35, 20)	middle frontal gyrus (dorsolateral prefrontal cortex)
BA46	(45, 35, 20)	middle frontal gyrus (dorsolateral prefrontal cortex)

MNI coordinates of the centroid voxels representative for the BAs and the respective underlying brain regions constituting the significant network obtained from the NBS analyses.

<https://doi.org/10.1371/journal.pone.0202600.t001>

and pars triangularis and the corresponding homologues in the right hemisphere (see Fig 1). The SIs did not show increased functional connectivity in any other of the tested frequency ranges compared to the controls nor did the controls show significant hyperconnectivities compared to the SIs. Neither did we observe a significant group difference in network connectivity measures during the resting state eyes closed condition.

Brain-behavior relationships

Correlation analyses between the mean connectivity value of the network obtained from the NBS analysis and autobiographical data of language characteristics within the SIs did not reveal significant results (mean connectivity value with YoE: $r = -0.006$, $p = 0.98$; training hours per week: $r = 0.074$, $p = 0.79$; training hours during life: $r = 0.202$, $p = 0.45$; AoC: $r = 0.430$, $p = 0.10$).

Discussion

In the present source-based EEG study, we revealed an increased functional connectivity matrix in SIs compared to multilingual control participants in the lower alpha frequency band during the eyes open period of the resting state condition. The obtained network was restricted to ventral and dorsal frontal regions across both hemispheres. In particular, SIs demonstrated




Fig 1. Significant network of the NBS analysis between the SIs and the controls in the lower alpha (8 Hz– 10 Hz) frequency band is shown in the sagittal, horizontal and coronal views (the blue spheres represent the location of the centroid voxels of the 84 BAs that entered the NBS analysis). SIs show increased functional connectivity (depicted in red) in a network containing BA9, BA44, 45 and 46 (enlarged blue spheres) spanning both hemispheres ($p = 0.046$, FWE-corrected). A = anterior, L = left hemisphere, R = right hemisphere.

<https://doi.org/10.1371/journal.pone.0202600.g001>

increased bilateral functional connectivity between the DLPFC and the ventral part of the prefrontal cortex, including pars opercularis, pars triangularis, and their right-sided homologues.

The frontal lobe regulates higher-order cognitive control mechanisms allowing successful performance of complex behavior as well as adapting to environmental changes [59]. Executive functions, an umbrella term for prefrontal processes, include among others attention, working memory, inhibition, initiation, self-monitoring and regulation [60]. Simultaneous interpreting as well as the control of multiple languages as in subjects who grew up bilingually brings along an intense training of higher-order cognitive functions to manage the selection of the proper language, switching between languages or translating and interpreting [3, 9, 61]. In fact, studies in the context of bilingualisms showed that bilingually grown up subjects outperform monolinguals in a variety of linguistic and non-linguistic tasks relying on executive functions such as inhibition [62, 63], working memory [64], attention [62, 65], and set shifting [66].

The sparse literature on neural activation patterns during the process of simultaneous interpreting reports a frontal activation increase in Broca's area when translating into the native language (L2 to L1) and an increase of activation in the left DLPFC when translating into the non-native language [6]. Comparably, literature of bilinguals also report an increased activity in the DLPFC when translating single words compared to word repetition [67]. In particular, an enhanced involvement of the DLPFC [68–70] as well as Broca's area [14] was observed during the process of language switching, and thus, constituting at least a subpart of an attentional network needed for language retrieval. However, to date it is still under debate which executive functions underlie language control. Both the DLPFC as well as Broca's area are discussed in terms of language-specific control processes and in being involved in general language-unrelated cognitive functions [71–74]. For example, described as subserving various functions [13, 75, 76], the Broca's area has gathered considerable attention in the domain of speech and language processing [77–82]. Based on a cytoarchitectonical view, Broca's area can be subdivided in a caudal (pars opercularis, BA 44) and a rostral (pars triangularis, BA 45) part [83]. From a functional perspective, Broca's area has previously been proposed to contribute to the unification of semantic and syntactic information at the sentence level. In fact, the pars opercularis contributes to planning of articulation, word retrieval memory, syntactic, and phonological processing [20, 78, 81]. Accordingly, pars triangularis has repeatedly been observed during diverse cognitive tasks such as language-related switching mechanisms, attention and inhibition [13–15, 84–86] as well as verbal working memory on both single word [87] and sentence processing [87–89]. Furthermore, an involvement of the pars triangularis has been particularly responsive to syntactic and semantic processes [11, 78, 79, 90–92].

Otherwise, the DLPFC plays an important role in prosodic processing [12, 76, 93] and fundamentally contributes to executive functions [94]. Furthermore, common prefrontal activation patterns in the DLPFC have previously been described during working memory processes and episodic memory processes [95–97]. Evidence from a microstructural point of view argues for a generous amount of inhibitory interneurons in the prefrontal cortex [98], thereby providing an ideal neural basis for cognitive control of language selection by actively suppressing the unwanted output in a face of competition [61]. Thus, the neural underpinnings of bilingualism emphasize the specific role of the prefrontal cortex for mediating inhibition-driven control of language switching, i.e., choosing the proper output language and eventual error correction [15, 70].

Based on 1) such nesting relations between linguistic and cognitive functions in the prefrontal cortex, 2) previous neuroimaging studies indicating structural differences between professional SIs and multilingual control participants in Broca's area and in its right-sided homologue that correlated with the amount of training [3], and 3) consistent activation patterns in Broca's area during simultaneous language translation across participants [99], we propose that simultaneous interpreting training may facilitate the neural communication

between spatially close areas situated in the ventral and dorsal parts of the prefrontal cortex. Such a processing mode seems to be particularly cost-effective in that it facilitates the local intertwining between linguistic and cognitive functions within local prefrontal circuits. This perspective is not only in line with hierarchical models of speech and language processing [11, 73, 92] but also with a previous diffusion tensor imaging (DTI) study that revealed anatomical differences in the anterior part of the corpus callosum between professional SIs and multilingual control participants [4]. Furthermore, the clustering of prefrontal areas into a densely interconnected functional matrix has previously been shown to be related to enhanced memory performance and cognitive flexibility [100]. This leads to suggest that dynamic training-related functional network reconfigurations constitute a fundamental neural principle underlying executive functions and cognition.

Increased alpha power has consistently been reported to be related to enhanced internally directed attention [101, 102] and to be negatively correlated to neural activity in the dorsal attention network [103]. These findings might lead to suggest that alpha activity is associated with cortical inhibitory processes tuning the brain into an optimized excitatory-inhibitory processing mode by suppressing the activity in neural networks that might interfere with proper processing of an ongoing task [28, 104]. In the context of language processing, mechanisms of language control or switching were described via actively blocking the non-target language by not accessing the mental lexicon of the irrelevant one [15, 105]. Disentangling the functional role of the alpha sub-bands, modulations of lower alpha power were shown to be driven as a function of tonic attentional demands [28, 106] and to be linked to better performance in a working memory task, whereas the upper alpha band seems to be related to stimulus-related cognitive processes [107].

Although we are aware that simultaneous interpreting requires various aspects of cognitive control mechanisms, two facts might support a slight dominance of inhibitory processes in SIs—at least the ones which are detectable during resting state. First, although oscillatory activity in the alpha frequency range is discussed within the frame of attention or working memory, it is also observed to function as a sub-process supporting memory and attention in a phase-dependent inhibitory manner [29]. Second, if working memory would play an important role in SIs we would have also expected to find an altered functional connectivity in the theta frequency range, the frequency band of many neural models of working memory, which was however not the case in our study [107, 108]. A more precise interpretation of our results is not possible at the moment lacking cognitive and behavioral data to infer clearly defined brain-behavior relations.

The differences of our findings compared to the ones of the fMRI resting state study by Becker and colleagues on SIs, consecutive interpreters and translators might be explained by the fact that acquisition modalities as well as MNI coordinates of the ROIs differed in the two studies [21]. Furthermore, Becker and colleagues did not have a multilingual control group without any interpreting experience, which makes it difficult comparing both studies directly.

We did not find significant correlations between the mean functional connectivity of our obtained network and language characteristics within the SI group. This does however not necessarily disqualify our findings as being expertise-related. It rather leads us to assume that either the variable mediating functional connectivity in SIs has not been properly acquired or that the obtained network displays a domain-specific predisposition rather than an experience- or training-related neural activity pattern. Therefore, future studies should for example apply approved objective language tests instead of self-assessed proficiency levels in a longitudinal approach.

The fact that the two groups did only differ in connectivity measures during the eyes open condition is not surprising. Previous research has shown that oscillatory activity during rest

with eyes open is not directly comparable with neural activation patterns during an eyes closed resting state period. Specifically, the dominant alpha oscillations observable during the eyes closed period are desynchronized or blocked when opening the eyes [109]. Besides this alpha dominance mediated by the reticular activation system [110, 111], alpha power also shows a positive relation with the arousal level, i.e. the current energetic level of the organisms [112] of the recorded subject across different resting state conditions. In this context, Barry and colleagues showed that the eyes open period was associated with an increased arousal level and a global reduction in alpha power compared to the eyes closed condition [113]. This conditional effect of alpha frequency power across the two resting state conditions recorded in the present study might have led to the fact that the expertise-related resting state network obtained during the eyes open period in the oscillatory alpha range was superimposed by the physiologically-driven change in alpha power during the eyes closed period. Differences across these two conditions have also been reported with regard to connectivity measures and in a graph-theoretical framework in both the alpha and theta frequency bands [114, 115]. In addition, as SIs interpret with eyes open, it is not surprising that a training-related network alteration appears in the more comparable condition.

To summarize, as being the first EEG study on the EEG source level, our results make a further contribution to the influence of language expertise on the intrinsic imprint in network activity patterns during a task-free resting state condition. Comparable to previous resting state studies in the context of expertise research, we found an altered connectivity in networks constituted of typical domain-related nodes. Our results support the idea of a training-related functional network configuration in SIs facilitating challenging language tasks by intertwining general cognitive and language-related processing. However, since we examined resting state network activity, the interpretation still remains speculative and therefore, further research in this area is needed.

Limitations

One limitation is that only female subjects participated in the present study. This is owed to the fact that it was nearly impossible to find enough male SIs. Besides diverse differences in resting state electroencephalographic activity and brain anatomy [116] between male and females [117], sexual dimorphisms of brain regions involved in speech processing have been observed [118–121]. In particular, language processing in females seems to be more bilaterally represented than in males who show a stronger left-dominant lateralization [122, 123]. Thus, future studies comparing network characteristics during a task-free condition between male and female (controlled for menstrual phase) SIs are necessary to draw straightforward conclusions. Furthermore, including a control group of other language experts, such as consecutive interpreters or translators might help to further unravel cognitive control mechanisms in SIs and the influence of language proficiency on connectivity patterns.

Supporting information

S1 File. English and German version of the in-house questionnaire on language skills.
(PDF)

S2 File. English and German version of the in-house questionnaire on musicality.
(PDF)

S3 File. Minimal data set.
(XLSX)

S1 Table. Autobiography of language characteristics. The age of acquisition (AoA), the years of experience (YoE), and self-reported linguistic skills (LS) in speaking (s) and writing (w) are listed in parenthesis for every learned language. Mother tongue is printed in bold.

SI = simultaneous interpreter, C = control subject.

(PDF)

S2 Table. Interpreting directions of the simultaneous interpreters. The symbol ‘↔’ means that SIs can translate into both languages, whereas ‘→’ means that SIs can only work in one direction. SI = simultaneous interpreter.

(PDF)

Acknowledgments

We thank all simultaneous interpreters and control subjects for participating in our study.

Author Contributions

Conceptualization: Carina Klein, Stefan Elmer, Lutz Jäncke.

Data curation: Carina Klein, Silvana Iris Metz.

Formal analysis: Carina Klein.

Funding acquisition: Lutz Jäncke.

Investigation: Carina Klein, Lutz Jäncke.

Methodology: Carina Klein, Lutz Jäncke.

Project administration: Carina Klein, Silvana Iris Metz, Stefan Elmer, Lutz Jäncke.

Resources: Carina Klein, Lutz Jäncke.

Software: Carina Klein.

Supervision: Carina Klein, Stefan Elmer, Lutz Jäncke.

Validation: Carina Klein.

Visualization: Carina Klein.

Writing – original draft: Carina Klein.

Writing – review & editing: Carina Klein, Silvana Iris Metz, Stefan Elmer, Lutz Jäncke.

References

1. Christoffels I, Degroot A, Kroll J. Memory and language skills in simultaneous interpreters: The role of expertise and language proficiency. *Journal of Memory and Language*. 2006; 54(3):324–45.
2. Elmer S, Meyer M, Jancke L. Simultaneous interpreters as a model for neuronal adaptation in the domain of language processing. *Brain Res*. 2010; 1317:147–56. <https://doi.org/10.1016/j.brainres.2009.12.052> PMID: 20051239
3. Elmer S, Hanggi J, Jancke L. Processing demands upon cognitive, linguistic, and articulatory functions promote grey matter plasticity in the adult multilingual brain: Insights from simultaneous interpreters. *Cortex*. 2014; 54:179–89. <https://doi.org/10.1016/j.cortex.2014.02.014> PMID: 24699036
4. Elmer S, Hanggi J, Meyer M, Jancke L. Differential language expertise related to white matter architecture in regions subserving sensory-motor coupling, articulation, and interhemispheric transfer. *Hum Brain Mapp*. 2011; 32(12):2064–74. <https://doi.org/10.1002/hbm.21169> PMID: 21162044
5. Hervais-Adelman A, Moser-Mercer B, Murray MM, Golestani N. Cortical thickness increases after simultaneous interpretation training. *Neuropsychologia*. 2017; 98:212–219. <https://doi.org/10.1016/j.neuropsychologia.2017.01.008> PMID: 28077311

6. Rinne JO, Tommola J, Laine M, Krause BJ, Schmidt D, Kaasinen V, et al. The translating brain: Cerebral activation patterns during simultaneous interpreting. *Neurosci Letters*. 2000; 294(2):85–8.
7. Hervais-Adelman A, Moser-Mercer B, Golestani N. Brain functional plasticity associated with the emergence of expertise in extreme language control. *Neuroimage*. 2015; 114:264–74. <https://doi.org/10.1016/j.neuroimage.2015.03.072> PMID: [25869858](#)
8. Abutalebi J. Neural aspects of second language representation and language control. *Acta Psychol (Amst)*. 2008; 128(3):466–78.
9. Abutalebi J, Della Rosa PA, Green DW, Hernandez M, Scifo P, Keim R, et al. Bilingualism tunes the anterior cingulate cortex for conflict monitoring. *Cereb Cortex*. 2012; 22(9):2076–86. <https://doi.org/10.1093/cercor/bhr287> PMID: [22038906](#)
10. Luk G, Green DW, Abutalebi J, Grady C. Cognitive control for language switching in bilinguals: A quantitative meta-analysis of functional neuroimaging studies. *Lang Cogn Process*. 2012; 27(10):1479–88.
11. Friederici AD, Gierhan SM. The language network. *Curr Opin Neurobiol*. 2013; 23(2):250–4. <https://doi.org/10.1016/j.conb.2012.10.002> PMID: [23146876](#)
12. Sammler D, Grosbras MH, Anwender A, Bestelmeyer PE, Belin P. Dorsal and Ventral Pathways for Prosody. *Curr Biol*. 2015; 25(23):3079–85. <https://doi.org/10.1016/j.cub.2015.10.009> PMID: [26549262](#)
13. Fedorenko E, Duncan J, Kanwisher N. Language-selective and domain-general regions lie side by side within Broca's area. *Curr Biol*. 2012; 22(21):2059–62. <https://doi.org/10.1016/j.cub.2012.09.011> PMID: [23063434](#)
14. Price CJ, Green DW, von Studnitz R. A functional imaging study of translation and language switching. *Brain*. 1999; 122:2221–35. PMID: [10581218](#)
15. Rodriguez-Fornells A, Rotte M, Heinze HJ, Nosselt T, Munte TF. Brain potential and functional MRI evidence for how to handle two languages with one brain. *Nature*. 2002; 415(6875):1026–9. <https://doi.org/10.1038/4151026a> PMID: [11875570](#)
16. Golestani N, Zatorre RJ. Learning new sounds of speech: Reallocation of neural substrates. *Neuroimage*. 2004; 21(2):494–506. <https://doi.org/10.1016/j.neuroimage.2003.09.071> PMID: [14980552](#)
17. Turkeltaub PE, Coslett HB. Localization of sublexical speech perception components. *Brain Lang*. 2010; 114(1):1–15. <https://doi.org/10.1016/j.bandl.2010.03.008> PMID: [20413149](#)
18. Ackermann H, Riecker A. The contribution(s) of the insula to speech production: A review of the clinical and functional imaging literature. *Brain Struct Funct*. 2010; 214(5–6):419–33. <https://doi.org/10.1007/s00429-010-0257-x> PMID: [20512374](#)
19. Dronkers NF. A new brain region for coordinating speech articulation. *Nature*. 1996; 384(6605):159–61. <https://doi.org/10.1038/384159a0> PMID: [8906789](#)
20. Eickhoff SB, Heim S, Zilles K, Amunts K. A systems perspective on the effective connectivity of overt speech production. *Philos Trans A Math Phys Eng Sci*. 2009; 367(1896):2399–421. <https://doi.org/10.1098/rsta.2008.0287> PMID: [19414462](#)
21. Becker M, Schubert T, Strobach T, Gallinat J, Kuhn S. Simultaneous interpreters vs. professional multilingual controls: Group differences in cognitive control as well as brain structure and function. *Neuroimage*. 2016; 134:250–60. <https://doi.org/10.1016/j.neuroimage.2016.03.079> PMID: [27085505](#)
22. Klein C, Liem F, Hanggi J, Elmer S, Jancke L. The "silent" imprint of musical training. *Hum Brain Mapp*. 2016; 37(2):536–46. <https://doi.org/10.1002/hbm.23045> PMID: [26538421](#)
23. Duan X, Long Z, Chen H, Liang D, Qiu L, Huang X, et al. Functional organization of intrinsic connectivity networks in Chinese-chess experts. *Brain Res*. 2014; 1558:33–43. <https://doi.org/10.1016/j.brainres.2014.02.033> PMID: [24565926](#)
24. Di X, Zhu S, Jin H, Wang P, Ye Z, Zhou K, et al. Altered resting brain function and structure in professional badminton players. *Brain Connect*. 2012; 2(4):225–33. <https://doi.org/10.1089/brain.2011.0050> PMID: [22840241](#)
25. Burzynska AZ, Finc K, Taylor BK, Knecht AM, Kramer AF. The dancing brain: Structural and functional signatures of expert dance training. *Front Hum Neurosci*. 2017; 11:566. <https://doi.org/10.3389/fnhum.2017.00566> PMID: [29230170](#)
26. Ward LM. Synchronous neural oscillations and cognitive processes. *Trends in Cognitive Sciences*. 2003; 7(12):553–9. PMID: [14643372](#)
27. Moreno S, Wodniecka Z, Tays W, Alain C, Bialystok E. Inhibitory Control in Bilinguals and Musicians: Event related potential (ERP) evidence for experience-specific effects. *Plos One*. 2014; 9(4).
28. Klimesch W. EEG alpha and theta oscillations reflect cognitive and memory performance: A review and analysis. *Brain Res Brain Res Rev*. 1999; 29(2–3):169–95. PMID: [10209231](#)

29. Klimesch W. Alpha-band oscillations, attention, and controlled access to stored information. *Trends Cogn Sci*. 2012; 16(12):606–17. <https://doi.org/10.1016/j.tics.2012.10.007> PMID: [23141428](#)
30. Annett M. A classification of hand preference by association analysis. *Br J Psychol*. 1970; 61:303–21. PMID: [5457503](#)
31. Elmer S, Klein C, Kuhn J, Liem F, Meyer M, Jancke L. Music and language expertise influence the categorization of speech and musical sounds: Behavioral and electrophysiological measurements. *J Cogn Neurosci*. 2014; 26(10):2356–69. https://doi.org/10.1162/jocn_a_00632 PMID: [24702451](#)
32. Elmer S, Kuhn J. Functional connectivity in the left dorsal stream facilitates simultaneous language translation: An EEG study. *Front Hum Neurosci*. 2016; 10:60. <https://doi.org/10.3389/fnhum.2016.00060> PMID: [26924976](#)
33. Gordon EE. Manual for the advanced measures of music education. Chicago: G.I.A. Publications, Inc.; 1989.
34. Elmer S, Meyer M, Jancke L. Neurofunctional and behavioral correlates of phonetic and temporal categorization in musically trained and untrained subjects. *Cereb Cortex*. 2012; 22(3):650–8. <https://doi.org/10.1093/cercor/bhr142> PMID: [21680844](#)
35. Elmer S, Hanggi J, Meyer M, Jancke L. Increased cortical surface area of the left planum temporale in musicians facilitates the categorization of phonetic and temporal speech sounds. *Cortex*. 2013; 49(10):2812–21. <https://doi.org/10.1016/j.cortex.2013.03.007> PMID: [23628644](#)
36. Lehl S, Gallwitz A, Blaha V, Fischer B. Theorie und Messung der geistigen Leistungsfähigkeit mit dem Kurztest KAI. Vless Ebersberg. 1991; 3. Aufl.
37. Lehl S. Mehrfachwahl-Wortschatz-Intelligenztest (MWT-B). Perimed, Erlangen. 1977.
38. Klein C, Diaz Hernandez L, Koenig T, Kottlow M, Elmer S, Jancke L. The influence of pre-stimulus EEG activity on reaction time during a verbal Sternberg task is related to musical expertise. *Brain Topogr*. 2016; 29(1):67–81. <https://doi.org/10.1007/s10548-015-0433-7> PMID: [25929715](#)
39. Jung TP, Makeig S, Humphries C, Lee TW, Mckeown MJ, Iragui V, et al. Removing electroencephalographic artifacts by blind source separation. *Psychophysiology*. 2000; 37:163–78. PMID: [10731767](#)
40. Friston KJ. Functional and effective connectivity: A review. *Brain connectivity*. 2011; 1(1):13–36. <https://doi.org/10.1089/brain.2011.0008> PMID: [22432952](#)
41. Babiloni C, Binetti G, Cassetta E, Cerboneschi D, Dal Forno G, Del Percio C, et al. Mapping distributed sources of cortical rhythms in mild Alzheimer's disease: A multicentric EEG study. *Neuroimage*. 2004; 22(1):57–67. <https://doi.org/10.1016/j.neuroimage.2003.09.028> PMID: [15109997](#)
42. Lehmann D, Faber PL, Gianotti LR, Kochi K, Pascual-Marqui RD. Coherence and phase locking in the scalp EEG and between LORETA model sources, and microstates as putative mechanisms of brain temporo-spatial functional organization. *Journal of Physiology (Paris)*. 2006; 99:29–36.
43. Mulert C, Gallinat J, Pascual-Marqui RD, Dorn H, DFrück K, Schlattmann P, et al. Reduced event-related current density in the anterior cingulate cortex in schizophrenia. *Neuroimage*. 2001; 13:589–600. <https://doi.org/10.1006/nimg.2000.0727> PMID: [11305888](#)
44. Pizzagalli D, Lehmann D, Koenig T, Regard M, Pascual-Marqui RD. Face-elicited ERPs and affective attitude: brain electric microstate and tomography analyses. *Clin Neurophysiol*. 2000; 111:521–31. PMID: [10699416](#)
45. Hirota T, Yagyu T, Pascual-Marqui RD, Saio N, Kinoshita T. Spatial structure of brain electric fields during intermittent photic stimulation. *Neuropsychobiology*. 2001; 44:108–12. <https://doi.org/10.1159/000054927> PMID: [11490182](#)
46. Seeck M, Lazeyras F, Michel CM, Blanke O, Gericke CA, Ives J, et al. Non-invasive epileptic focus localization using EEG-triggered functional MRI and electromagnetic tomography. *Electroencephalogr Clin Neurophysiol*. 1998; 106(6):508–12. PMID: [9741750](#)
47. Worrell GA, Lagerlund TD, Sharbough FW, Brinkmann BH, Busacker NE, Cicora KM, et al. Localization of the epileptic focus by low-resolution electromagnetic tomography in patients with a lesion demonstrated by MRI. *Brain Topogr*. 2000; 12(4):273–82. PMID: [10912735](#)
48. Dierks T, Jelic V, Pascual-Marqui RD, Wahlund LO, Julin P, Linden DEJ, et al. Spatial pattern of cerebral glucose metabolism (PET) correlates with localization of intracerebral EEG-generators in Alzheimer's disease. *Clin Neurophysiol*. 2000; 111:1817–24. PMID: [11018498](#)
49. Pascual-Marqui RD. Coherence and phase synchronization: generalization to pairs of multivariate time series, and removal of zero-lag contributions. *arXiv:07061776v3 [statME]*. 2007.
50. Brodmann K. Vergleichende Lokalisationslehre der Grosshirnrinde in ihren Prinzipien dargestellt auf Grund des Zellenbaues. Barth, Leipzig; 1909.

51. Pascual-Marqui RD, Lehmann D, Koukkou M, Kochi K, Anderer P, Saletu B, et al. Assessing interactions in the brain with exact low-resolution electromagnetic tomography. *Philos Trans A Math Phys Eng Sci*. 2011; 369(1952):3768–84. <https://doi.org/10.1098/rsta.2011.0081> PMID: 21893527
52. Langer N, von Bastian CC, Wirz H, Oberauer K, Jancke L. The effects of working memory training on functional brain network efficiency. *Cortex*. 2013; 49(9):2424–38. <https://doi.org/10.1016/j.cortex.2013.01.008> PMID: 23489778
53. Jancke L, Langer N. A strong parietal hub in the small-world network of coloured-hearing synaesthetes during resting state EEG. *J Neuropsychol*. 2011; 5(2):178–202. <https://doi.org/10.1111/j.1748-6653.2011.02004.x> PMID: 21923785
54. De Vico Fallani F, Maglione A, Babiloni F, Mattia D, Astolfi L, Vecchiato G, et al. Cortical network analysis in patients affected by schizophrenia. *Brain Topogr*. 2010; 23(2):214–20. <https://doi.org/10.1007/s10548-010-0133-2> PMID: 20094766
55. Pascual-Marqui RD. Instantaneous and lagged measurements of linear and nonlinear dependence between groups of multivariate time series: frequency decomposition. *ArXiv*. 2007;arXiv:0711.1455 [stat.ME](<http://arxiv.org/abs/0711.1455>).
56. Zalesky A, Fornito A, Bullmore ET. Network-based statistic: identifying differences in brain networks. *Neuroimage*. 2010; 53(4):1197–207. <https://doi.org/10.1016/j.neuroimage.2010.06.041> PMID: 20600983
57. Nichols TE, Holmes AP. Nonparametric permutation tests for functional neuroimaging experiments: A primer with examples. *Hum Brain Mapp*. 2002; 15:1–25. PMID: 11747097
58. Xia M, Wang J, He Y. BrainNet Viewer: a network visualization tool for human brain connectomics. *PLoS One*. 2013; 8(7):e68910. <https://doi.org/10.1371/journal.pone.0068910> PMID: 23861951
59. Otero TM, Barker LA. The Frontal Lobes and Executive Functioning. *Handbook of Executive Functioning*. 2014. p. 29–44.
60. Goldstein S, Naglieri JA, Princiotta D, Otero TM. Introduction: a history of executive functioning. In Goldstein S. & Naglieri J.A. (Eds.). *Handbook of Executive Functioning*. 2013; New York, NY: Springer.
61. Abutalebi J, Green D. Bilingual language production: The neurocognition of language representation and control. *Journal of Neurolinguistics*. 2007; 20(3):242–75.
62. Bialystok E, Craik FI, Klein R, Viswanathan M. Bilingualism, aging, and cognitive control: Evidence from the Simon task. *Psychol Aging*. 2004; 19(2):290–303. <https://doi.org/10.1037/0882-7974.19.2.290> PMID: 15222822
63. Festman J, Rodriguez-Fornells A, Münte TF. Individual differences in control of language interference in late bilinguals are mainly related to general executive abilities. *Behav Brain Funct*. 2010; 6(5).
64. Morales J, Calvo A, Bialystok E. Working memory development in monolingual and bilingual children. *J Exp Child Psychol*. 2013; 114(2):187–202. <https://doi.org/10.1016/j.jecp.2012.09.002> PMID: 23059128
65. Costa A, Hernandez M, Sebastian-Galles N. Bilingualism aids conflict resolution: Evidence from the ANT task. *Cognition*. 2008; 106(1):59–86. <https://doi.org/10.1016/j.cognition.2006.12.013> PMID: 17275801
66. Festman J, Münte TF. Cognitive control in Russian-german bilinguals. *Front Psychol*. 2012; 3:115. <https://doi.org/10.3389/fpsyg.2012.00115> PMID: 22529831
67. Klein D, Milner B, Zatorre RJ, Meyer E, Evans AC. The neural substrates underlying word generation: A bilingual functional-imaging study. *Proc Natl Acad Sci U S A*. 1995; 92(7):2899–903. PMID: 7708745
68. Hernandez AE, Dapretto M, Mazziotta J, Bookheimer S. Language switching and language representation in Spanish-English bilinguals: An fMRI study. *Neuroimage*. 2001; 14(2):510–20. <https://doi.org/10.1006/nimg.2001.0810> PMID: 11467923
69. Hernandez AE, Martinez A, Kohnert K. In search of the language switch: An fMRI study of picture naming in Spanish-English bilinguals. *Brain Lang*. 2000; 73(3):421–31. <https://doi.org/10.1006/brln.1999.2278> PMID: 10860563
70. Hernandez AE. Language switching in the bilingual brain: what's next? *Brain Lang*. 2009; 109(2–3):133–40. <https://doi.org/10.1016/j.bandl.2008.12.005> PMID: 19250662
71. Vigneau M, Beaucois V, Herve PY, Duffau H, Crivello F, Houde O, et al. Meta-analyzing left hemisphere language areas: Phonology, semantics, and sentence processing. *Neuroimage*. 2006; 30(4):1414–32. <https://doi.org/10.1016/j.neuroimage.2005.11.002> PMID: 16413796
72. Vigneau M, Beaucois V, Herve PY, Jobard G, Petit L, Crivello F, et al. What is right-hemisphere contribution to phonological, lexico-semantic, and sentence processing? Insights from a meta-analysis.

- Neuroimage. 2011; 54(1):577–93. <https://doi.org/10.1016/j.neuroimage.2010.07.036> PMID: [20656040](#)
73. Hagoort P. On Broca, brain, and binding: A new framework. *Trends Cogn Sci*. 2005; 9(9):416–23. <https://doi.org/10.1016/j.tics.2005.07.004> PMID: [16054419](#)
74. Hagoort P. Nodes and networks in the neural architecture for language: Broca's region and beyond. *Curr Opin Neurobiol*. 2014; 28:136–41. <https://doi.org/10.1016/j.conb.2014.07.013> PMID: [25062474](#)
75. Fedorenko E. The role of domain-general cognitive control in language comprehension. *Front Psychol*. 2014; 5:335. <https://doi.org/10.3389/fpsyg.2014.00335> PMID: [24803909](#)
76. Friederici AD. The brain basis of language processing: From structure to function. *Physiol Rev*. 2011; 91(4):1357–92. <https://doi.org/10.1152/physrev.00006.2011> PMID: [22013214](#)
77. Bookheimer S. Functional MRI of language: New approaches to understanding the cortical organization of semantic processing. *Annu Rev Neurosci*. 2002; 25:151–88. <https://doi.org/10.1146/annurev.neuro.25.112701.142946> PMID: [12052907](#)
78. Friederici AD. Broca's area and the ventral premotor cortex: Functional differentiation and specificity. *Cortex*. 2006; 42(4):472–5. PMID: [16881252](#)
79. Fiebach CJ, Schlesewsky M, Lohmann G, von Cramon DY, Friederici AD. Revisiting the role of Broca's area in sentence processing: Syntactic integration versus syntactic working memory. *Hum Brain Mapp*. 2005; 24(2):79–91. <https://doi.org/10.1002/hbm.20070> PMID: [15455462](#)
80. Price CJ. The anatomy of language: Contributions from functional neuroimaging. *J Anat*. 2000; 197(3):335–59.
81. Price CJ. A review and synthesis of the first 20 years of PET and fMRI studies of heard speech, spoken language and reading. *Neuroimage*. 2012; 62(2):816–47. <https://doi.org/10.1016/j.neuroimage.2012.04.062> PMID: [22584224](#)
82. Meyer ME, Jäncke L. Involvement of the left and right frontal operculum in speech and nonspeech perception and production. in *Broca's Region*. 2006; Oxford University Press:218–41.
83. Amunts K, Schleicher A, Bürgel U, Mohlberg H, Uylings HB, Zilles K. Broca's region revisited: Cytoarchitecture and intersubject variability. *J Comp Neurol*. 1999; 412(2):319–41. PMID: [10441759](#)
84. Fuster JM. The prefrontal cortex—an update: Time is of the essence. *Neuron*. 2001; 30:319–33. PMID: [11394996](#)
85. Santi A, Grodzinsky Y. Broca's area and sentence comprehension: A relationship parasitic on dependency, displacement or predictability? *Neuropsychologia*. 2012; 50(5):821–32. <https://doi.org/10.1016/j.neuropsychologia.2012.01.017> PMID: [22285904](#)
86. Rodriguez-Fornells A, de Diego-Balaguer R, Münte TF. Executive control in bilingual language processing. *Language Learning*. 2006; 56(s1):133–90.
87. Chein JM, Fissell K, Jacobs S, Fiez JA. Functional heterogeneity within Broca's area during verbal working memory. *Physiol Behav*. 2002; 77(4–5):635–9. PMID: [12527011](#)
88. Rogalsky C, Hickok G. The role of Broca's area in sentence comprehension. *J Cogn Neurosci*. 2011; 23(7):1664–80. <https://doi.org/10.1162/jocn.2010.21530> PMID: [20617890](#)
89. Caplan D, Waters GS. Verbal working memory and sentence comprehension. *Behav Brain Sci*. 1999; 22(1):77–94. PMID: [11301522](#)
90. Thompson CK, Tait ME, Ballard KJ, Fix SC. Agrammatic aphasic subject's comprehension of subject and object extracted Wh questions. *Brain Lang*. 1999; 67(3):169–87. <https://doi.org/10.1006/brln.1999.2052> PMID: [10210629](#)
91. Tyler LK, Marslen-Wilson WD, Randall B, Wright P, Devereux BJ, Zhuang J, et al. Left inferior frontal cortex and syntax: Function, structure and behaviour in patients with left hemisphere damage. *Brain*. 2011; 134(Pt 2):415–31. <https://doi.org/10.1093/brain/awq369> PMID: [21278407](#)
92. Bornkessel-Schlesewsky I, Schlesewsky M. Reconciling time, space and function: A new dorsal-ventral stream model of sentence comprehension. *Brain Lang*. 2013; 125(1):60–76. <https://doi.org/10.1016/j.bandl.2013.01.010> PMID: [23454075](#)
93. Strelnikov KN, Vorobyev VA, Chernigovskaya TV, Medvedev SV. Prosodic clues to syntactic processing—a PET and ERP study. *Neuroimage*. 2006; 29(4):1127–34. <https://doi.org/10.1016/j.neuroimage.2005.08.021> PMID: [16188459](#)
94. Duncan J, Owen AM. Common regions of the human frontal lobe recruited by diverse cognitive demands. *Trends Neurosci*. 2000; 23(10):475–83. PMID: [11006464](#)
95. Blumenfeld RS, Parks CM, Yonelinas AP, Ranganath C. Putting the pieces together: The role of dorsolateral prefrontal cortex in relational memory encoding. *J Cogn Neurosci*. 2011; 23(1):257–65. <https://doi.org/10.1162/jocn.2010.21459> PMID: [20146616](#)

96. Curtis CE, D'Esposito M. Persistent activity in the prefrontal cortex during working memory. *Trends Cogn Sci*. 2003; 7(9):415–23. PMID: [12963473](#)
97. Balconi M. Dorsolateral prefrontal cortex, working memory and episodic memory processes: insight through transcranial magnetic stimulation techniques. *Neurosci Bull*. 2013; 29(3):381–9. <https://doi.org/10.1007/s12264-013-1309-z> PMID: [23385388](#)
98. Miller EK, Erickson CA, Desimone R. Neural mechanisms of visual working memory in prefrontal cortex of the macaque. *J Neurosci*. 1996; 16(16):5154–67. PMID: [8756444](#)
99. Elmer S. Broca pars triangularis constitutes a "hub" of the language-control network during simultaneous language translation. *Front Hum Neurosci*. 2016; 10:491. <https://doi.org/10.3389/fnhum.2016.00491> PMID: [27746729](#)
100. Braun U, Schafer A, Walter H, Erk S, Romanczuk-Seiferth N, Haddad L, et al. Dynamic reconfiguration of frontal brain networks during executive cognition in humans. *Proc Natl Acad Sci U S A*. 2015; 112(37):11678–83. <https://doi.org/10.1073/pnas.1422487112> PMID: [26324898](#)
101. Palva S, Palva JM. New vistas for alpha-frequency band oscillations. *Trends Neurosci*. 2007; 30(4):150–8. <https://doi.org/10.1016/j.tins.2007.02.001> PMID: [17307258](#)
102. Cooper NR, Croft RJ, Dominey SJ, Burgess AP, Gruzelier JH. Paradox lost? Exploring the role of alpha oscillations during externally vs. internally directed attention and the implications for idling and inhibition hypotheses. *Int J Psychophysiol*. 2003; 47(1):65–74. PMID: [12543447](#)
103. Laufs H, Kleinschmidt A, Beyerle A, Eger E, Salek-Haddadi A, Preibisch C, et al. EEG-correlated fMRI of human alpha activity. *Neuroimage*. 2003; 19(4):1463–76. PMID: [12948703](#)
104. Sadaghiani S, Kleinschmidt A. Functional interactions between intrinsic brain activity and behavior. *Neuroimage*. 2013; 80:379–86. <https://doi.org/10.1016/j.neuroimage.2013.04.100> PMID: [23643921](#)
105. de Groot AM, Delmaar P, Lupker SJ. The processing of interlexical homographs in translation recognition and lexical decision: Support for non-selective access to bilingual memory. *Q J Exp Psychol A*. 2000; 53(2):397–428.
106. Klimesch W, Doppelmayr M, Russegger H, Pachinger T, Schwaiger J. Induced alpha band power changes in the human EEG and attention. *Neurosci Lett*. 1998; 244(2):73–6. PMID: [9572588](#)
107. Klimesch W. EEG-alpha rhythms and memory processes. *Int J Psychophysiol*. 1997; 26(1–3):319–40. PMID: [9203012](#)
108. Lisman JE, Jensen O. The theta-gamma neural code. *Neuron*. 2013; 77(6):1002–16. <https://doi.org/10.1016/j.neuron.2013.03.007> PMID: [23522038](#)
109. Berger H. Über das Elektrenkephalogramm des Menschen. *Archly für Psychiatrie*. 1933; 98:231–54.
110. Gale A, Coles MGH, Boyd E. Variation in visual input and the occipital EEG: II. *Psychon Science*. 1971; 23:99–100.
111. Mayhew SD, Ostwald D, Porcaro C, Bagshaw AP. Spontaneous EEG alpha oscillation interacts with positive and negative BOLD responses in the visual-auditory cortices and default-mode network. *Neuroimage*. 2013; 76:362–72. <https://doi.org/10.1016/j.neuroimage.2013.02.070> PMID: [23507378](#)
112. Pribram KH, McGuinness D. Arousal, activation, and effort in the control of attention. *Psychological Review*. 1975; 82(2):116–49. PMID: [1096213](#)
113. Barry RJ, Clarke AR, Johnstone SJ, Magee CA, Rushby JA. EEG differences between eyes-closed and eyes-open resting conditions. *Clin Neurophysiol*. 2007; 118(12):2765–73. <https://doi.org/10.1016/j.clinph.2007.07.028> PMID: [17911042](#)
114. Jin SH, Jeong W, Lee DS, Jeon BS, Chung CK. Preserved high-centrality hubs but efficient network reorganization during eyes-open state compared with eyes-closed resting state: an MEG study. *J Neurophysiol*. 2014; 111(7):1455–65. <https://doi.org/10.1152/jn.00585.2013> PMID: [24431400](#)
115. Tan B, Kong X, Yang P, Jin Z, Li L. The difference of brain functional connectivity between eyes-closed and eyes-open using graph theoretical analysis. *Comput Math Methods Med*. 2013; 2013:976365. <https://doi.org/10.1155/2013/976365> PMID: [23690886](#)
116. Sowell ER, Peterson BS, Kan E, Woods RP, Yoshii J, Bansal R, et al. Sex differences in cortical thickness mapped in 176 healthy individuals between 7 and 87 years of age. *Cereb Cortex*. 2007; 17(7):1550–60. <https://doi.org/10.1093/cercor/bhl066> PMID: [16945978](#)
117. Polunina AG, Lefterova NP. Gender differences in resting state electroencephalography characteristics. *Current Trends in Neurology*. 2012; 6:51–60.
118. Luders E, Narr KL, Thompson PM, Rex DE, Woods RP, Deluca H, et al. Gender effects on cortical thickness and the influence of scaling. *Hum Brain Mapp*. 2006; 27(4):314–24. <https://doi.org/10.1002/hbm.20187> PMID: [16124013](#)

119. Schlaepfer TE, Harris GJ, Tien AY, Peng L, Lee S, Pearlson GD. Structural differences in the cerebral cortex of healthy female and male subjects: a magnetic resonance imaging study. *Psychiatric Res: Neuroimaging*. 1995; 61:129–35. PMID: [8545497](#)
120. Kurth F, Jancke L, Luders E. Sexual dimorphism of Broca's region: More gray matter in female brains in Brodmann areas 44 and 45. *J Neurosci Res*. 2017; 95(1–2):626–32. <https://doi.org/10.1002/jnr.23898> PMID: [27870461](#)
121. Wallentin M. Putative sex differences in verbal abilities and language cortex: a critical review. *Brain Lang*. 2009; 108(3):175–83. <https://doi.org/10.1016/j.bandl.2008.07.001> PMID: [18722007](#)
122. Shaywitz BA, Shaywitz SE, Pugh KR, Constable RT, Skudlarski P, Fulbright RK, et al. Sex differences in the functional organization of the brain for language. *Nature*. 1995; 373(6515):607–9. <https://doi.org/10.1038/373607a0> PMID: [7854416](#)
123. Vikingstad EM, George KP, Johnson AF, Cao Y. Cortical language lateralization in right handed normal subjects using functional magnetic resonance imaging. *J Neurol Sci*. 2000; 175(1):17–27. PMID: [10785252](#)